



***Unit-Dose Non-Aerosol Spray
20% Benzocaine Oral Anesthetic***

Directions For Use

1. Tear at notch to open pouch. Remove HurriCaine ONE from pouch.
2. To unlock, turn clockwise.
3. Grasp HurriCaine ONE firmly by placing your index and middle fingers on the finger grip and positioning your thumb against the base of the plunger.
4. Position patient with mouth open. Aim spray tip toward the back of patient's throat.
5. Apply firm pressure to the plunger by pushing your thumb forward to initiate the spray action. Instruct patient to swallow 1-2 times. Anesthesia is accomplished within 30 seconds and lasts approximately 15 minutes.

Active ingredient: Benzocaine 20%

Purpose: Oral Anesthetic

Uses: For the temporary relief of occasional minor irritation and pain associated with sore mouth and throat

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- Pale, gray, or blue colored skin (cyanosis)
- Headache
- Rapid heart rate
- Shortness of breath
- Dizziness or lightheadedness
- Fatigue or lack of energy

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Do not use

- if foil pouch has been previously opened
- for teething
- in children under 2 years of age

When using this product avoid contact with eyes

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions: Do not exceed recommended dosage

Adults and children 2 years of age and older: use up to 4 times daily or as directed by a dentist or doctor

Children under 12 years of age: should be supervised in the use of the product

Children under 2 years of age: do not use

Other information: store at 15-30 °C (59-86 °F)

Inactive ingredients: flavor, polyethylene glycol, sodium saccharin

Questions or comments?

1-800-238-8542

M-F: 8:00 a.m. – 4:30 p.m. ET