


## DIRECTIONS


### Step 1

#### Remove cap



### STEP 2

#### Affix syringe tip



### STEP 3

#### Apply product to area



- Do not exceed recommended dosage
- For adults and children 12 years and older
- Attach a disposable tip onto syringe
- Prior to scaling and root planing, apply a small amount of gel along the gingival margin and directly into periodontal pockets on the buccal and lingual surfaces; apply to one quadrant at a time. Anesthesia will take place within 30 seconds and last approximately 15 minutes.
- Discard syringe and tip after use on each patient
- HurriCaine® may be applied up to 4 times daily

### Additional uses

For direct application during scaling:

- Prior to periodontal scaling and root planing, take the syringe that contains HurriCaine Topical Anesthetic and dispense it into a dappen dish.
- Dip the scaler end into the dappen dish to coat the tip for a direct application of anesthetic during treatment. Reapply to scaler as necessary.

### For irrigation to site specific application:

- Attach a disposable tip onto syringe
- Irrigate along the gingival margin or into the periodontal pocket where anesthesia is needed

## Ordering Information

HurriCaine Topical Anesthetic Gel - Original Wild Cherry - Syringe.....NDC #0283-0520-29

HurriCaine Topical Anesthetic Gel - Piña Colada - Syringe .....NDC #0283-1820-29

**Beutlich**  
**PHARMACEUTICALS LLC**  
 Since 1954

To place an order or for more information, please contact your dental supplier or call us at 1-800-238-8542, M-F: 8:00 a.m. - 4:30 p.m. ET. HurriCaine is a registered trademark of Beutlich® Pharmaceuticals, LLC.  
 www.beutlich.com

HSIAF 897 0125

## Drug Facts

### Active ingredient

Benzocaine 20%.....Oral Anesthetic

### Purpose

**Uses** for the temporary relief of occasional minor irritation and pain, associated with:
 

- canker sores
- sore mouth and throat
- minor injury of the mouth and gums
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances

### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:
 

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

**Do not use**

- if cap is missing or damaged
- for teething
- in children under 2 years of age

**When using this product** avoid contact with eyes

**Keep out of reach of children.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

### Directions

do not exceed recommended dosage  
 adults and children apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.

children under 12 years of age	should be supervised in the use of the product
children under 2 years of age	do not use

### Other information

store at 20-25 °C (68-77 °F)

### Inactive ingredients

flavor, polyethylene glycol, sodium saccharin

### Questions or comments?

1-800-238-8542

M-F: 8:00 a.m. - 4:30 p.m. ET