




## DIRECTIONS




1. Peel back film



2. Pop-up cotton swab



3. Secure gel tray



4. Dip cotton swab in gel and apply product to area to anesthetize

## Drug Facts

### Active ingredient

Benzocaine 20%.....Oral Anesthetic

### Purpose

**Uses** for the temporary relief of occasional minor irritation and pain, associated with

- canker sores
- sore mouth and throat
- minor injury of the mouth and gums
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances

### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

## Drug Facts (continued)

### Warnings (continued)

**Do not use**

- if product film is punctured or damaged
- for teething
- in children under 2 years of age

**When using this product** avoid contact with eyes

**Keep out of reach of children.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

### Directions

do not exceed recommended dosage adults and children apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.

children under 12 years of age	should be supervised in the use of the product
children under 2 years of age	do not use

### Other information

store at 20-25 °C (68-77 °F)

### Inactive ingredients

flavor, polyethylene glycol, sodium saccharin

### Questions or comments?

1-800-238-8542

M-F: 8:00 a.m. – 4:30 p.m. ET

## Ordering Information

HurriCaine Unit Dose Topical Anesthetic Gel - Original Wild Cherry - 60 ct. .... NDC 0283-0871-59  
HurriCaine Unit Dose Topical Anesthetic Gel - Watermelon - 60 ct. .... NDC 0283-0293-59  
HurriCaine Unit Dose Topical Anesthetic Gel - Strawberry - 60 ct. .... NDC 0283-1016-59