



Single-Use 4% Lidocaine Spray

Directions For Use

1. Tear at notch to open pouch. Remove HurriCaine ONE from pouch.
2. To unlock, turn clockwise.
3. Grasp HurriCaine ONE Lidocaine firmly by placing your index and middle fingers on the finger grip and positioning your thumb against the base of the plunger.
4. Apply firm pressure to the plunger by pushing your thumb forward to initiate the spray action. Numbing effect is achieved within 5 minutes and lasts 10+ minutes after application.

Active ingredients: Lidocaine 4% and Benzalkonium chloride 0.13%

Purpose: Topical Analgesic and First Aid Antiseptic

Uses: For temporary relief of pain. First aid to help protect against infection in minor cuts, scrapes, and burns.

Warnings:

- For external use only. Do not use in the eyes or apply over large areas of the body, particularly over raw surfaces or blistered areas. In case of deep or puncture wounds, animal bites, or serious burns, consult a doctor.
- If condition worsens, or if symptoms persist more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor. Do not use longer than 1 week unless directed by a doctor.
- Flammable, Keep away from fire or flame.

Do not use

- If foil pouch has been previously opened
- In children under 2 years of age

When using this product avoid contact with eyes.

Keep out of reach of children. If accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions: Do not exceed recommended dosage

Adults and children 2 years of age and older: Clean the affected area. Spray a small amount of this product on the area 1 to 3 times daily. May be covered with a sterile bandage. If bandaged, let dry first.

Children under 2 years of age: do not use, consult a physician.

Other information: store at 15-30°C (59-86°F)

Inactive ingredients: Ethanol, Polyethylene glycol

Questions or comments?

1.800.238.8542

M-F 7:30 am – 4:00 pm ET